ADHD EDUCATION | 3 MIN READ

Understanding Executive Function

How ADHD affects the brain's ability to plan, focus, remember, and get things done.

Have you ever walked into a room and forgotten why? Started homework and ended up doing something else? Everyone experiences this sometimes. For people with ADHD, these difficulties can happen often enough to disrupt everyday life.

What is executive function?

Imagine your brain is a busy airport. The air traffic controller helps planes take off, land, and avoid collisions. Executive functions are like that controller: they help you organize, control impulses, manage emotions, remember instructions, and finish tasks. The prefrontal cortex works with other brain regions to support these skills.

Six important executive-function skills

Working memory: Hold information long enough to use it, such as remembering three instructions.

Attention control: Focus on an important task despite distractions.

Planning and organization: Break big projects into steps and keep track of materials.

Impulse control: Pause and think before speaking or acting.

Flexible thinking: Adjust when plans change.

Task initiation: Get started even when a task feels boring or difficult.

How does ADHD affect executive function?

ADHD involves differences in networks that support attention, motivation, and self-regulation. Dopamine helps regulate reward, learning, and goal-directed behavior. Norepinephrine helps regulate alertness and attention. Both are neurotransmitters: chemical messengers between nerve cells. ADHD is not explained by a single simple chemical shortage. A child may understand homework but struggle to begin; an adult may know about an appointment but still lose track of time. This is not automatically laziness or low intelligence.

How can medication help?

ADHD medicines may make it easier to regulate attention and behavior. They do not automatically teach organization or time-management skills.

Medication group	Simplified action
Methylphenidate (Ritalin, Concerta)	Reduces dopamine and norepinephrine reuptake.
Amphetamines (Adderall, Vyvanse)	Increase catecholamine signaling via release and transporter effects.
Atomoxetine (Strattera)	Primarily blocks norepinephrine reuptake.
Guanfacine ER (Intuniv)	Stimulates alpha-2A receptors to support prefrontal networks.
Clonidine ER (Kapvay)	Stimulates alpha-2 adrenergic receptors; affects attention and arousal.

Five ways to strengthen these skills

Use reminders: Calendars, alarms, and checklists reduce memory demands.

Break tasks into steps: Start with one small, specific action.

Create routines: Predictable study, sleep, and preparation times help.

Reduce distractions: Use a quiet workspace and limit unnecessary alerts.

Celebrate progress: Recognize small improvements, not only perfect results.

When should you seek help?

Consider an ADHD evaluation when persistent problems with focus, organization, impulse control, or completing tasks significantly affect school, work, relationships, or daily responsibilities. An evaluation also considers anxiety, depression, sleep problems, and learning difficulties.

Key message: ADHD does not mean someone cannot succeed. Supportive environments, practical skills, and sometimes clinical treatment can help people use their abilities effectively.

References

Diamond A. Executive functions. Annual Review of Psychology. 2013;64:135–168. doi:10.1146/annurev-psych-113011-143750.

National Institute of Mental Health. Attention-Deficit/Hyperactivity Disorder. https://www.nimh.nih.gov/health/topics/attention-deficit-hyperactivity-disorder-adhd

Wolraich ML, et al. Clinical practice guideline for ADHD in children and adolescents. Pediatrics. 2019;144(4):e20192528. doi:10.1542/peds.2019-2528.

Arnsten AFT. The emerging neurobiology of ADHD: the key role of the prefrontal association cortex. Journal of Pediatrics. 2009;154(5):I-S43. doi:10.1016/j.jpeds.2009.01.018.

24-7 Shalom Psychiatry PLLC | General education only; not a substitute for individualized assessment or treatment.